

MEMBER HANDBOOK

IMMUNIZATION (SHOT) SCHEDULE FOR CHILDREN

Immunizations (shots) help prevent serious illness. This record will help keep track when your child is immunized. If your child did not get their shots at the age shown, they still need to get that shot. Talk to your PCP about your child's immunization (shots). Children must have their immunizations (shots) to enter school.

| IMMUNIZATION RECORD | | |
|----------------------|--|---------------|
| Age | Shot (Immunization) | Date Received |
| Birth | НерВ | |
| 2 Months | DTaP, Hib, IPV, PCV, RV, HepB | |
| 4 Months | DTaP, Hib, IPV, PCV, RV | |
| 6 Months | DTaP, Hib, IPV, PCV, RV, HepB | |
| 12-15 Months | Hib, PCV, MMR, Varicella, HepA* | |
| 15-18 Months | DTaP** | |
| 19-23 Months | HepA* | |
| 4-6 years | DTaP, IPV, MMR, Varicella | |
| 7-10 years Catch-Up | Tdap, HepB, IPV, MMR, Varicella, HepA | |
| 11-12 years | Tdap, MenACWY (1 dose), HPV (2 doses)*** | |
| 11-12 years Catch-Up | HepB, IPV, MMR, Varicella, HepA | |
| 13-18 years Catch-Up | Tdap, MenACWY (1 dose, Booster at 16), HPV, HepB, IPV, MMR, Varicella, HepA | |
| 16-18 years | MenB**** | |
| 19-20 years | HPV (3 doses) *****, MMR*****, Tdap*****, Varicella***** | |
| Every Year | Influenza (after 6 months) | |

*If the birth dose of HepB is given, the 4 month dose may be omitted.

**Can be given as early as 12 months, if there are six months since third dose.

***a 3-shot series is needed for those with weakened immune systems and those age 15 or older.

****Can be given sooner for those that have certain health conditions that would put them at increased risk for serious diseases.

*****Recommended unless your healthcare provider tells you that you do not need it or should not get

it.



| IMMUNIZATION RECORD | | |
|-------------------------|---|---------------|
| Age | Shot (Immunization) | Date Received |
| Birth | НерВ | |
| 2 months | DTaP, Hib, IPV, PCV, RV, HepB | |
| 4 months | DTaP, Hib, IPV, PCV, RV | |
| 6 months | DTaP, Hib, IPV, PCV, RV, HepB | |
| 12-15 months | Hib, PCV, MMR, Varicella, HepA* | |
| 15-18 months | DTaP** | |
| 19-23 months | HepA* | |
| 4-6 years | DTaP, IPV, MMR, Varicella | |
| 7-10 years Catch-Up | Tdap, HepB, IPV, MMR, Varicella, HepA | |
| 11-12 years | Tdap, MenACWY (1 dose), HPV (2 doses)*** | |
| 11-12 years Catch-Up | HepB, IPV, MMR, Varicella , HepA | |
| 13-18 years Catch-Up | Tdap, MenACWY (1 dose, Booster at 16), MenB (16-18 years)*****, HPV (2 doses)****, HepB, IPV, MMR, Varicella, HepA | |
| 16-18 years | MenB**** | |
| 19-20 years | HPV – (2-3 doses)****, MMR****, Tdap****, Varicella**** | |
| Every year | Influenza (after 6 months) | |

* The fist dose of HepA vaccine should be given between 12 months and 23

months of age. The second dose should be given 6 months after the fist dose.

**Can be given as early as 12 months,

if there are six months since third dose.

***A 3rd shot series is needed for those

with weakened immune systems and those who start the series at 15 years or older.

****Recommended unless your health-

care provider tells you that you cannot safely receive it or that you do not need it. HPV doses depending on age of initialvaccination or condition.

*****Ages 16-18 who are not at increased

risk may receive after discussing with their healthcare provider.